arXiv:1605.02478v1 [math.NT] 9 May 2016

SQUARE-FULL POLYNOMIALS IN SHORT INTERVALS AND IN ARITHMETIC PROGRESSIONS

E. RODITTY-GERSHON

Abstract. We study the variance of sums of the indicator function of square-full polynomials in both arithmetic progressions and short intervals. Our work is in the context of the ring $\mathbb{F}_q[T]$ of polynomials over a finite field $\mathbb{F}_q$ of q elements, in the limit $q \to \infty$. We use a recent equidistribution result due to N. Katz to express these variances in terms of triple matrix integrals over the unitary group, and evaluate them.

Contents

1. Introduction 1
2. Square-full Polynomials 3
2.1. Arithmetic progressions 3
2.2. Short intervals 4
3. Dirichlet characters and Katz's equidistribution results 5
3.1. Dirichlet characters 5
3.2. Dirichlet L-functions 8
3.3. Katz's equidistribution results 8
4. The variance in arithmetic progressions 9
4.1. The mean value 9
4.2. The case of small n 10
4.3. A formula for the variance 10
4.4. The quadratic character and the cubic character 11
4.5. Average of the sum $\mathcal{M}(n; \alpha_2\chi)$ 11
4.6. Proof of Theorem 2.1 13
5. The variance over short intervals 14
5.1. The mean value 14
5.2. An expression for the variance 15
Acknowledgements 18
References 18

1. Introduction

A positive integer n is called a square-full number if $p^2|n$ for every prime factor p of n. Denote by α_2 the indicator function of square-full numbers,

Date: May 10, 2016.

i.e.

$$\alpha_2(n) = \begin{cases} 1 & \text{if } n \text{ is square-full} \\ 0 & \text{otherwise} \end{cases} \tag{1.1}$$

Let $\mathcal{A}(x)$ be the number of square full integers not exceeding x. In 1935, Erdös and Szekeres [5] proved

$$\mathcal{A}(x) = \frac{\zeta(3/2)}{\zeta(3)} x^{1/2} + O_\epsilon(x^{1/3+\epsilon}). \tag{1.2}$$

Bateman and Grosswald [1] improved this result in 1958. They obtained

$$\mathcal{A}(x) = \frac{\zeta(3/2)}{\zeta(3)} x^{1/2} + \frac{\zeta(2/3)}{\zeta(2)} x^{1/3} + O(x^{1/6} \cdot \mathrm{e}^{-\mathrm{c}(\log^3 \mathrm{x}/\log\log \mathrm{x})^{\frac{1}{5}}}) \tag{1.3}$$

where c is a positive absolute constant. They also made the observation that any improvment of the exponent $\frac{1}{6}$ would imply that $\zeta(s) \neq 0$ for $\Re s > 1-\delta$ ($\delta > 0$). There is a very long history of studies and conditional improvements of the error term in the above formula (see [2],[3],[4],[15],[19],[20],[21],[22]).

It follows from (1.3) that for intervals of length $H > x^{2/3+\epsilon}$ we have

$$\sum_{x \leq n \leq x+H} \alpha_2(n) \sim \frac{\zeta(3/2)}{2\zeta(3)} \cdot H/x^{1/2}. \tag{1.4}$$

Various authors used exponential sum techniques to reduce the lower bound on H for which this asymptotic is valid. Heath-Brown [7] proved it with for $H > x^{\eta+\epsilon}$ with $\eta = 0.6318\ldots$. Liu [16] proved it with $\eta = 0.6308\ldots$. Filaseta and Trifonov[6] found a simpler approach, using real instead of complex analysis, and obtained the exponent $\eta = 0.6282\ldots$. In [8], Huxley and Trifonov improve this to $H \geq \frac{1}{\epsilon} x^{5/8} (\log x)^{5/16}$.

Concerning the distribution of square full numbers in arithmetic progressions, the most recent result is due to Munsch [18]. By evaluating character sums, he showed that

$$\sum_{\substack{n \leq x \\ n = a \bmod q}} \alpha_2(n) \sim \frac{\zeta(3/2)}{\zeta(3)} \frac{A_{a,q}}{q} x^{1/2} + \frac{\zeta(2/3)}{\zeta(2)} \frac{B_{a,q}}{q} x^{1/3} \tag{1.5}$$

where

$$A_{a,q} = \prod_{p|q} (1 - \frac{1}{p^3})^{-1} \sum_{\chi \in X_2} \chi(a) \frac{L(3/2, \chi)}{\zeta(3/2)}$$

and

$$B_{a,q} = \prod_{p|q} (1 - \frac{1}{p^2})^{-1} \sum_{\chi \in X_3} \bar{\chi}(a) \frac{L(2/3, \chi)}{\zeta(2/3)}$$

with X_2 and X_3 being the set of all quadratic and cubic characters mod q respectively and $L(s, \chi)$ is the L-function attached χ.

The goal of this paper is to study the fluctuations of the analogous sums in the function field settings. Namely, we study the variance of the sum of α_2 in arithmetic progressions or in short intervals, in the context of the ring

$\mathbb{F}_q[T]$ of polynomials over a finite field $\mathbb{F}_q$ of q elements, in the limit $q \to \infty$. In our setting we succeed in giving definitive answers in both cases.

Our approach involves converting the problem to one about the correlation of zeros of a certain family of L-functions, and then using an equidistribution result of Katz which holds in the limit $q \to \infty$.

2. Square-full Polynomials

Let $\mathbb{F}_q$ be a finite field of an odd cardinality q, and let M_n be the set of all monic polynomials of degree n with coefficients in $\mathbb{F}_q$. In analogy to numbers, we say that $f \in \mathcal{M}_n$ is a square-full polynomial if for every polynomial $P \in \mathcal{M}_n$ that divides f, P^2 also divides f. We denote by α_2 the indicator function of square-full polynomials, i.e.

$$\alpha_2(f) = \begin{cases} 1 & \text{if } f \text{ is square-full} \\ 0 & \text{otherwise} \end{cases} \tag{2.1}$$

The mean value of α_2 over all monic polynomials is defined to be

$$\langle \alpha_2 \rangle_n := \frac{1}{q^n} \sum_{f \in \mathcal{M}_n} \alpha_2(f). \tag{2.2}$$

The generating function for the number of monic square-full polynomials of degree n, i.e. $\sum_{f \in M_n} \alpha_2(f)$ is

$$\sum_{n=0}^{\infty} \sum_{f \in M_n} \alpha_2(f) u^n = \frac{Z(u^2)Z(u^3)}{Z(u^6)} \tag{2.3}$$

where $Z(u)$ is the zeta function of $\mathbb{F}_q[T]$ (also set $\zeta_q := Z(q^{-s})$) , given by the following product over prime polynomials in $\mathbb{F}_q[T]$

$$Z(u) = \prod_P (1 - u^{\deg P})^{-1} = \frac{1}{1 - qu} \tag{2.4}$$

By expanding an comparing coefficients we have for $n > 6$

$$\sum_{f \in M_n} \alpha_2(f) = \frac{q^{n/2}}{\zeta_q(3)} \sum_{\substack{j = n \bmod 2 \\ 0 \le j \le \lfloor \frac{n}{3} \rfloor - 2}} q^{\frac{-j}{2}} + q^{\lfloor \frac{n - \lfloor n/3 \rfloor}{2} \rfloor} \tag{2.5}$$

therefore in the limit of $q \to \infty$ we get

$$\langle \alpha_2 \rangle_n \sim q^{\lfloor n/2 \rfloor - n}. \tag{2.6}$$

2.1. **Arithmetic progressions.**

Let $Q \in \mathbb{F}_q[T]$ be a squarefree polynomial of a positive degree. The sum of α_2 over all monic polynomials of degree n lying in the arithmetic progressions $f = A \bmod Q$ is

$$S_{\alpha_2;n;Q}(A) := \sum_{\substack{f \in \mathcal{M}_n \\ f = A \bmod Q}} \alpha_2(f) \tag{2.7}$$

The average of this sum $S_{\alpha_2;n;Q}(A)$ when we vary A over residue classes coprime to Q is

$$\langle S_{\alpha_2;n;Q}\rangle = \frac{1}{\Phi(Q)} \sum_{\substack{f\in\mathcal{M}_n \\ (f,Q)=1}} \alpha_2(f) \tag{2.8}$$

where $\Phi(Q)$ is the number of invertible residues modulo Q.

In section 4 we will consider the variance of $S_{\alpha_2;n;Q}$ which is defined to be the average of the squared difference between $S_{\alpha_2;n;Q}$ and its mean value

$$\mathrm{Var}(S_{\alpha_2;n;Q}) = \frac{1}{\Phi(Q)} \sum_{\substack{A \bmod Q \\ (A,Q)=1}} |S_{\alpha_2;n;Q} - \langle S_{\alpha_2;n;Q}\rangle|^2. \tag{2.9}$$

proving the following Theorem:

Theorem 2.1. *Let Q be a prime polynomial and set $N := \deg Q - 1$, then in the limit $q \to \infty$ the following holds*
for $N \le n \le 2N$

$$\mathrm{Var}(S_{\alpha_2;n;Q}) \sim \frac{q^{\lfloor \frac{n}{2} \rfloor}}{\Phi(Q)}$$

for $2N < n$ even

$$\mathrm{Var}(S_{\alpha_2;n;Q}) \sim \frac{q^n}{\Phi(Q)^2}$$

for $2N < n$ odd

$$\mathrm{Var}(S_{\alpha_2;n;Q}) \sim \frac{q^{n-3}}{\Phi(Q)^2} \cdot |\sum_{j=1}^{\deg Q-1} \alpha_j(\chi_2)|^2 = O(\frac{q^{n-2}}{\Phi(Q)^2})$$

where χ_2 is the quadratic character mod Q and $\alpha_j(\chi_2)$ are the inverse roots of the L-function associated to χ_2.

Note that the restriction that Q is a prime is for simplicity reasons only.

2.2. **Short intervals.** A "short interval" in $\mathbb{F}_q[x]$ is a set of the form

$$I(A;h) = \{f : ||f - A|| \le q^h\} \tag{2.10}$$

where $A \in \mathcal{M}_n$ and $0 \le h \le n-2$. the norm is

$$||f|| := \#\mathbb{F}_q[t]/(f) = q^{\deg f}\ . \tag{2.11}$$

The cardinality of such a short interval is

$$\#I(A;h) = q^{h+1} := H\ . \tag{2.12}$$

We define for $1 \le h < n$ and $A \in \mathcal{M}_n$

$$\mathcal{N}_{\alpha_2;h}(A) := \sum_{f\in I(A;h)} \alpha_2(f) \tag{2.13}$$

to be the number of square-full polynomials in the interval $I(A;h)$.

The mean value of $\mathcal{N}_{\alpha_2;h}$ when we average over $A \in \mathcal{M}_n$ is

$$\langle \mathcal{N}_{\alpha_2;h} \rangle := \frac{1}{q^n} \sum_{A \in \mathcal{M}_n} \mathcal{N}_{\alpha_2;h}(A) \tag{2.14}$$

In section 5 we will compute the variance of $\mathcal{N}_{\alpha_2;h}$

$$\mathrm{Var}(\mathcal{N}_{\alpha_2;h}) := \frac{1}{q^n} \sum_{A \in \mathcal{M}_n} |\mathcal{N}_{\alpha_2;h}(A) - \langle \mathcal{N}_{\alpha_2;h} \rangle|^2 \tag{2.15}$$

proving the following Theorem:

Theorem 2.2. *Set $N := n - h - 2$, then in the limit $q \to \infty$ the following holds*
for $0 \leq n \leq 2N$

$$\mathrm{Var}(\mathcal{N}_{\alpha_2;h}) \sim \frac{H}{q^n} \cdot q^{\lfloor \frac{n}{2} \rfloor} \tag{2.16}$$

for $2N < n \leq 5N$

$$\mathrm{Var}(\mathcal{N}_{\alpha_2;h}) \sim \frac{H}{q^n} \cdot q^{\lfloor \frac{n+N}{3} \rfloor} \tag{2.17}$$

for $5N < n$

$$\mathrm{Var}(\mathcal{N}_{\alpha_2;h}) \sim \frac{H}{q^n} \cdot q^{\frac{n+N}{6}} \cdot q^{\frac{-\lambda_n}{6}} \tag{2.18}$$

where

$$\lambda_n = \begin{cases} 0 & n = 5N \mod 6 \\ 7 & n = 5N+1 \mod 6 \\ 6 & n = 5N+2 \mod 6 \\ 3 & n = 5N+3 \mod 6 \\ 4 & n = 5N+4 \mod 6 \\ 11 & n = 5N+5 \mod 6 \end{cases} \tag{2.19}$$

3. Dirichlet characters and Katz's equidistribution results

3.1. **Dirichlet characters.** Let $Q(T) \in \mathbb{F}_q[T]$ be a polynomial of positive degree. A Dirichlet character modulo Q is a homomorphism

$$\chi : (\mathbb{F}_q[T]/(Q))^\times \to \mathbb{C}^\times. \tag{3.1}$$

One can extend χ to $\mathbb{F}_q[T]$ by defining it to vanish on polynomials which are not coprime to Q. We denote by $\Gamma(Q)$ the group of all Dirichlet characters modulo Q. Note that $|\Gamma(Q)|$ is the Euler totient function $\Phi(Q)$. A Dirichlet character needs then to satisfy the following: $\chi(fg) = \chi(f)\chi(g)$ for all $f, g \in \mathbb{F}_q[T]$, $\chi(1) = 1$ and $\chi(f + hQ) = \chi(f)$ for all $f, h \in \mathbb{F}_q[T]$.

The orthogonality relations for Dirichlet characters are:

$$\frac{1}{\Phi(Q)}\sum_{\chi \bmod Q}\bar{\chi}(A)\chi(N)=\begin{cases}1 & N=A \bmod Q\\ 0 & \text{otherwise}\end{cases} \tag{3.2}$$

$$\frac{1}{\Phi(Q)}\sum_{A \bmod Q}\bar{\chi_1}(A)\chi_2(A)=\begin{cases}1 & \chi_1=\chi_2\\ 0 & \text{otherwise}\end{cases} \tag{3.3}$$

A character χ is called "even" if it acts trivially on the elements of $\mathbb{F}_q$, i.e. if $\chi(cf)=\chi(f)$ for all $0\neq c\in\mathbb{F}_q$. Therefore, the number of even characters is given by $\Phi^{ev}(Q)=\frac{\Phi(Q)}{q-1}$. For example there are q^{m-1} even character mod T^m. Out of this there are $O(q^{m-2})$ nonprimitive even characters mod T^m (see subsection 3.3 in [13]) . A character is called "odd" if it is not even.

A character χ is *primitive* if there is no proper divisor $Q'|Q$ such that $\chi(f)=1$ whenever f is co-prime to Q and $f=1 \bmod Q'$. Define the following:

$\Gamma_{prim}(Q)$-the set of all primitive characters mod Q, $|\Gamma_{prim}(Q)|:=\Phi_{prim}(Q)$.

$\Gamma^{ev}_{prim}(Q)$-the set of primitive even characters mod Q, $|\Gamma^{ev}_{prim}(Q)|:=\Phi^{ev}_{prim}(Q)$.

$\Gamma^{odd}_{prim}(Q)$-the set of primitive odd characters mod Q, $|\Gamma^{odd}_{prim}(Q)|:=\Phi^{odd}_{prim}(Q)$.

$\Gamma_{d-prim}(Q)$-the set of all characters χ mod Q such that χ^d is primitive for the fixed integer $d>0$, $|\Gamma_{d-prim}(Q)|:=\Phi_{d-prim}(Q)$. note that this is a subset of $\Gamma_{prim}(Q)$.

$\Gamma^{d-odd}_{d-prim}(Q)$-the set of all characters χ mod Q such that χ^d is primitive and odd for the fixed integer $d>0$, $|\Gamma^{d-odd}_{d-prim}(Q)|:=\Phi^{d-odd}_{d-prim}(Q)$.

Next, we will check the proportion of the set $\Gamma_{d-prim}(Q)$ in the the group of all characters mod Q ,$\Gamma(Q)$.

Lemma 3.1. *Let $Q\in\mathbb{F}_q[t]$ be a square-free polynomial, then in the limit of a large field size $q\to\infty$,*

$$\frac{\Phi_{d-prim}(Q)}{\Phi(Q)}=1+O(\frac{1}{q}) \tag{3.4}$$

Proof. Consider equation (3.21) in [13]. It asserts that as $q\to\infty$, almost all characters are **primitive** in the sense that

$$\frac{\Phi_{prim}(Q)}{\Phi(Q)}=1+O(\frac{1}{q}) \tag{3.5}$$

We claim, in the lemma above, that this also holds when $\Phi_{prim}(Q)$ is replaced by $\Phi_{d-prim}(Q)$. And indeed, if Q factors into k irreducible polynomials P_i

of degree d_i, $\deg Q := n = \sum_{i=1}^{k} d_i$ and $\Phi(Q) = \prod_{i=1}^{k}(q^{d_i} - 1)$. Therefore, $\Phi(Q) \sim q^n$ as $q \to \infty$. By the above combined with (3.5), we may conclude that the number of non primitive characters is $O(q^{n-1})$. Note that a character χ does not lie in $\Gamma_{d-prim}(Q)$ if χ^d does not lie in $\Gamma_{prim}(Q)$. In that case, χ^d lies in $\frac{\Gamma(Q)}{\Gamma_{prim}(Q)}$, and recall that $\#(\frac{\Gamma(Q)}{\Gamma_{prim}(Q)}) = O(q^{n-1})$. Now consider the map $\chi \mapsto \chi^d$. Its kernel is of cardinality $\#\{\chi|\chi^d = 1\}$. Since every χ is a product of characters χ_j of $\mathbb{F}_q[T]/(f_j)$ when $Q = \prod_{j=1}^{k} f_j$, $\deg(f_j) = d_j$ and $\sum_{i=1}^{k} d_i = n$, the following bound holds:

$$\#\{\chi|\chi^d = 1\} \leq d^n. \tag{3.6}$$

Since every character χ such that χ^d is not primitive can be written as a product of a non primitive character and an element of the kernel, we get that $\#(\frac{\Gamma(Q)}{\Gamma_{d-prim}(Q)})$ can be at most $d^n \times O(q^{n-1}) = O(q^{n-1})$, therefore we get (3.4). □

Note that equation (3.25) in [13] asserts that as $q \to \infty$, almost all characters are **primitive** and **odd** in the sense that

$$\frac{\Phi^{odd}_{prim}(Q)}{\Phi(Q)} = 1 + O(\frac{1}{q}) \tag{3.7}$$

Hence, exactly as before, we may also show that

Lemma 3.2. *Let $Q \in \mathbb{F}_q[t]$ be a square-free polynomial, then in the limit of a large field size $q \to \infty$,*

$$\frac{\Phi^{d-odd}_{d-prim}(Q)}{\Phi(Q)} = 1 + O(\frac{1}{q}) \tag{3.8}$$

Next, we will prove short lemma stating that under certain restrictions on the characteristic of the field, the primitivity of χ and χ^d is equivalent when χ is an even character mod T^m. This lemma will be useful later on in section 5.

Lemma 3.3. *Let d be an integer co-prime to $\Phi(Q)$. Then the map $\chi \mapsto \chi^d$ is an automorphism of the group of characters mod Q, i.e. an automorphism of $\Gamma(Q)$.*

Proof. The map is clearly an endomorphism since the group is abelian. Now, d is co-prime to the order of the group therefore there aren't any elements who's order dividing d and hence the kernel of the map is trivial. □

Lemma 3.4. *Let χ be an even Dirichlet character mod T^m, and let d be an integer s.t. $d < p$ when p is the characteristic of the field $\mathbb{F}_q$. Then χ is a primitive character if and only if χ^d is a primitive character.*

Proof. The order of the the subgroup of even characters mod T^m is $\Phi^{ev}(T^m) = q^{m-1}$. Taking $d < p$ when p is the characteristic of the field $\mathbb{F}_q$ gives d co-prime to $\Phi(T^m)$, in which case the above lemma applies. □

3.2. **Dirichlet L-functions.** The L-function associated to a dirichlet character $\chi \mod Q$ is defined as the following product over prime polynomials $P \in \mathbb{F}_q[T]$

$$L(u,\chi) = \prod_{P \nmid Q} (1 - \chi(P) u^{\deg P})^{-1} \tag{3.9}$$

The product is absolutely convergent for $|u| < 1/q$. For $\chi = \chi_0$ the trivial character mod Q

$$L(u,\chi_0) = Z(u) \prod_{P|Q} (1 - u^{\deg P}) \tag{3.10}$$

If $Q \in \mathbb{F}_q[T]$ is a polynomial of degree $\deg Q \geq 2$ and χ is a nontrivial character mod Q, then the L-function associated to χ i.e. $L(u,\chi)$ is a polynomial in u of degree $\deg Q - 1$. If χ is even then $L(u,\chi)$ has a trivial zero at $u = 1$. Now, we may factor $L(u,\chi)$ in terms of the inverse roots

$$L(u,\chi) = \prod_{j=1}^{\deg Q - 1} (1 - \alpha_j(\chi) u) \tag{3.11}$$

for which the Riemann Hypothesis, proved by Weil, asserts that for each (nonzero) inverse root, either $\alpha_j(\chi) = 1$ or

$$|\alpha_j(\chi)| = q^{1/2} \tag{3.12}$$

For a primitive odd character mod Q all the inverse roots have absolute value $q^{1/2}$. For a primitive even character mod Q all the inverse roots have absolute value $q^{1/2}$, except for the trivial zero at 1. Thus, we may write $\alpha_j(\chi) = q^{1/2} \mathrm{e}^{i\Theta_j}$, and the L-function (for a primitive character χ) is

(3.13)
$$L(u,\chi) = (1 - \lambda_\chi u)^{-1} \det(I - u q^{1/2} \Theta_\chi), \qquad \Theta_\chi = \mathrm{diag}(\mathrm{e}^{i\Theta_1}, \ldots, \mathrm{e}^{i\Theta_N}).$$

where $N = \deg Q - 1$ and $\lambda_\chi = 0$ for odd character χ. For even χ we have $N = \deg Q - 2$ and $\lambda_\chi = 1$. The unitary matrix $\Theta_\chi \in U(N)$ determines a unique conjugacy class which is called the unitarized Frobenius matrix of χ.

3.3. **Katz's equidistribution results.** The main ingredients in our results on the variance are equidistribution and independence results for the Frobenii Θ_χ due to N. Katz.

Theorem 3.5. [9] *Fix $m \geq 4$. The unitarized Frobenii Θ_χ for the family of even primitive characters $\mod T^{m+1}$ become equidistributed in the projective unitary group $PU(m-1)$ of size $m-1$, as q goes to infinity.*

Theorem 3.6. [11] *If $m \geq 5$ and in addition the characteristics of the fields $\mathbb{F}_q$ is bigger than 13, then the set of conjugacy classes $(\Theta_{\chi^2}, \Theta_{\chi^3}, \Theta_{\chi^6})$ become equidistributed in the space of conjugacy classes of the product $PU(m-1) \times PU(m-1) \times PU(m-1)$ as q goes to infinity.*

For odd characters, the corresponding equidistribution and independence results are

Theorem 3.7. [10] *Fix $m \geq 2$. Suppose we are given a sequence of finite fields $\mathbb{F}_q$ and squarefree polynomials $Q(T) \in \mathbb{F}_q[T]$ of degree m. As $q \to \infty$, the conjugacy classes Θ_χ with χ running over all primitive odd characters modulo Q, are uniformly distributed in the unitary group $U(m-1)$.*

Theorem 3.8. [12] *If in addition we restrict the characteristics of the fields $\mathbb{F}_q$ is bigger than 6, then the set of conjugacy classes $(\Theta_{\chi^2}, \Theta_{\chi^3}, \Theta_{\chi^6})$ with χ running over all characters such that χ^2, χ^3, χ^6 are primitive odd characters modulo Q,become equidistributed in the space of conjugacy classes of the product $U(m-1) \times U(m-1) \times U(m-1)$ as q goes to infinity.*

4. The variance in arithmetic progressions

4.1. **The mean value.** Given a polynomial $Q \in \mathbb{F}_q[T]$ the average of $S_{\alpha_2;n;Q}(A)$ when we vary A over residue classes co-prime to Q (see (2.7)) equals to

$$\langle S_{\alpha_2;n;Q} \rangle = \frac{1}{\Phi(Q)} \sum_{\substack{f \in \mathcal{M}_n \\ (f,Q)=1}} \alpha_2(f) = \frac{1}{\Phi(Q)} \sum_{f \in \mathcal{M}_n} \chi_0(f)\alpha_2(f) \tag{4.1}$$

To evaluate this consider the generating function

$$\begin{aligned}
\langle S_{\alpha_2;n;Q} \rangle &= \frac{1}{\Phi(Q)} \sum_{\substack{f \in \mathcal{M}_n \\ (f,Q)=1}} \alpha_2(f) \\
&= \frac{1}{\Phi(Q)} \cdot \sum_{f \in \mathcal{M}_n} \chi_0(f)\alpha_2(f) \\
&= \frac{1}{\Phi(Q)} \cdot \frac{L(u^2,\chi_0)L(u^3,\chi_0)}{L(u^6,\chi_0)} \qquad (4.2) \\
&= \frac{1}{\Phi(Q)} \cdot \frac{Z(u^2)Z(u^3)}{Z(u^6)} \prod_{P|Q} \left(\frac{(1-u^{2\deg P})(1-u^{3\deg P})}{(1-u^{6\deg P})}\right) \\
&= \frac{1}{\Phi(Q)} \cdot \frac{Z(u^2)Z(u^3)}{Z(u^6)} \prod_{P|Q} \left(\frac{(1-u^{2\deg P})}{(1+u^{3\deg P})}\right)
\end{aligned}$$

By expanding and comparing coefficients it is clear that the leading order coefficient in q (we are interested in $q \to \infty$) comes from $\frac{Z(u^2)Z(u^3)}{Z(u^6)}$. Therefor, by 2.6, we have

$$\langle S_{\alpha_2;n;Q} \rangle \sim \frac{q^{\lfloor n/2 \rfloor}}{\Phi(Q)}. \tag{4.3}$$

In the rest of this section we will evaluate the variance of $S_{\alpha_2;n;Q}$ i.e. the average of the squared difference between $S_{\alpha_2;n;Q}$ and its mean value.

4.2. **The case of small n.** See also subsection 4.2 in [14]. If $n < \deg Q$ then there is at most one polynomial $f \in \mathbb{F}_q[T]$ of degree n such that $f = A \mod Q$. In this case, when $q \to \infty$

$$Var(S_{\alpha_2;n;Q}) \sim \frac{q^n}{\Phi(Q)}\langle\alpha_2\rangle_n \tag{4.4}$$

Indeed, if $n < \deg Q$ we can use (4.3) to see that

$$|\langle S_{\alpha_2;n;Q}\rangle| \ll_n \frac{1}{q^{n/2}} \tag{4.5}$$

Hence

$$\begin{aligned} Var(S_{\alpha_2;n;Q}) =& \frac{1}{\Phi(Q)} \sum_{\substack{A \bmod Q \\ (A,Q)=1}} |S_{\alpha_2;n;Q}(A)|^2(1+O(\frac{1}{q^{n/2}})) \\ &= \frac{1}{\Phi(Q)} \sum_{\substack{f\in\mathcal{M}_n \\ (f,Q)=1}} |\alpha_2(f)|^2(1+O(\frac{1}{q^{n/2}})) \\ &\sim \frac{q^n}{\Phi(Q)}\langle\alpha_2\rangle_n \\ &\sim \frac{q^{\lfloor n/2\rfloor}}{\Phi(Q)} \end{aligned} \tag{4.6}$$

4.3. **A formula for the variance.** We present a formula for the variance of $S_{\alpha_2;n;Q}$ using Dirichlet characters [14, §4.1]. We start with the following expansion, using the first orthogonality relation for Dirichlet characters (see (3.2)) to pick out an arithmetic progression:

$$S_{\alpha_2,n,Q}(A) = \frac{1}{\Phi(Q)} \sum_{\substack{f\in\mathcal{M}_n \\ (f,Q)=1}} \alpha_2(f) + \frac{1}{\Phi(Q)} \sum_{\chi\neq\chi_0} \overline{\chi(A)}\mathcal{M}(n;\alpha_2\chi) \tag{4.7}$$

Where

$$\mathcal{M}(n;\alpha_2\chi) := \sum_{f\in M_n} \alpha_2(f)\chi(f)\ . \tag{4.8}$$

Note that the contribution of the trivial character equals to the average of $S_{\alpha_2;n;Q}(A)$. Therefore

$$S_{\alpha_2;n;Q} - \langle S_{\alpha_2;n;Q}\rangle = \frac{1}{\Phi(Q)} \sum_{\chi\neq\chi_0 \bmod Q} \overline{\chi}(A)\mathcal{M}(n;\alpha_2\chi) \tag{4.9}$$

Using the above and the second orthogonality relation for Dirichlet characters (see (3.3)), we have the following expression for the variance:

$$Var(S_{\alpha_2;n;Q}) = \langle|S_{\alpha_2;n;Q} - \langle S_{\alpha_2;n;Q}\rangle|^2\rangle = \frac{1}{\Phi(Q)^2} \sum_{\chi\neq\chi_0} |\mathcal{M}(n;\alpha_2\chi)|^2 \tag{4.10}$$

4.4. **The quadratic character and the cubic character.** Next, we evaluate $\mathcal{M}(n;\alpha_2\chi)$ for $\chi=\chi_2$ a quadratic character and for $\chi=\chi_3$ a cubic character. We assume here for simplicity that Q is prime polynomial. The sum $\mathcal{M}(n;\alpha_2\chi)$ given by (4.8), is the coefficient of u^n in the expansion of $\frac{L(u^2,\chi^2)L(u^3,\chi^3)}{L(u^6,\chi^6)}$. Thus for $\chi=\chi_2$, the generating function of $\mathcal{M}(n;\alpha_2\chi_2)$ has the following form

$$L(u^3,\chi_2)\cdot\frac{Z(u^2)(1-u^{2\deg Q})}{Z(u^6)(1-u^{6\deg Q})} \tag{4.11}$$

by expanding and comparing coefficients while bearing in mind the Riemann hypothesis (3.12), we get

$$\mathcal{M}(n;\alpha_2\chi_2)\sim\begin{cases} q^{\frac{n}{2}} & n\,\text{even}\\ -q^{\frac{n-3}{2}}\cdot\sum_{j=1}^{\deg Q-1}\alpha_j(\chi_2) & n\,\text{odd}\end{cases} \tag{4.12}$$

where $\alpha_j(\chi_2)$ are the inverse roots of $L(u^3,\chi_2)$.

For $\chi=\chi_3$, the generating function of $\mathcal{M}(n;\alpha_2\chi_3)$ has the following form

$$L(u^2,\chi_3^2)\cdot\frac{Z(u^3)(1-u^{3\deg Q})}{Z(u^6)(1-u^{6\deg Q})} \tag{4.13}$$

as before we get

$$\mathcal{M}(n;\alpha_2\chi_3)\sim\begin{cases} q^{\frac{n}{3}} & n=0 \mod 3\\ q^{\frac{n-4}{3}}\cdot\sum_{j,l=1}^{\deg Q-1}\alpha_j(\chi_3)\alpha_l(\chi_3) & n=1 \mod 3\\ -q^{\frac{n-2}{3}}\cdot\sum_{j=1}^{\deg Q-1}\alpha_j(\chi_3) & n=2 \mod 3\end{cases} \tag{4.14}$$

where $\alpha_j(\chi_3)$ are the inverse roots of $L(u^2,\chi_3)$.

4.5. **Average of the sum $\mathcal{M}(n;\alpha_2\chi)$.**

Lemma 4.1. *For a dirichlet character χ mod Q, such that χ^2,χ^3,χ^6 are odd and primitive*

$$\mathcal{M}(n;\alpha_2\chi)=\sum_{\substack{2j+3l+6k=n\\0\le j\le N\\0\le l\le N\\0\le k}} q^{\frac{j+k+l}{2}}\operatorname{tr}\Lambda_j(\Theta_{\chi^2})\operatorname{tr}\Lambda_l(\Theta_{\chi^3})\operatorname{tr}\operatorname{Sym}^k(\Theta_{\chi^6}) \tag{4.15}$$

Where $\Theta_\chi\in U(\deg Q-1)$ is the unitarized Frobenius matrix, Sym^n *is the symmetric $n-th$ power representation, and Λ_n is the exterior $n-th$ power representation.*
For $\chi\neq\chi_0,\chi_2,\chi_3$ mod Q not of the above, the following bound holds

$$|\mathcal{M}(n;\alpha_2\chi)|\ll_{\deg Q}\sum_{\substack{2j+3l+6k=n\\0\le j\le N\\0\le l\le N\\0\le k}} q^{\frac{j+k+l}{2}} \tag{4.16}$$

Proof. The sum $\mathcal{M}(n;\alpha_2\chi)$ given by (4.8), is the coefficient of u^n in the expansion of $\frac{L(u^2,\chi^2)L(u^3,\chi^3)}{L(u^6,\chi^6)}$. For an odd primitive characters χ mod Q, we use the Riemann Hypothesis (3.12) (Weils theorem) to write

$$L(u,\chi)=\det(I-uq^{\frac{1}{2}}\Theta_\chi)=\sum_{i=0}^{\deg Q-1}u^i q^{i/2}\operatorname{tr}\Lambda_i(\Theta_\chi) \tag{4.17}$$

and

$$\frac{1}{L(u,\chi)}=\frac{1}{\det(I-uq^{\frac{1}{2}}\Theta_\chi)}=\sum_{i=0}^{\infty}u^i q^{i/2}\operatorname{tr}\operatorname{Sym}^i(\Theta_\chi) \tag{4.18}$$

Abbreviate as follows:

$$\Lambda_i(\chi):=\operatorname{tr}\Lambda_i(\Theta_\chi),\ \operatorname{Sym}^i(\chi)=\operatorname{tr}\operatorname{Sym}^i(\Theta_\chi)$$

to have

$$\frac{L(u^2,\chi^2)L(u^3,\chi^3)}{L(u^6,\chi^6)}=\sum_{j=0}^{\deg Q-1}\sum_{l=0}^{\deg Q-1}\sum_{k=0}^{\infty}u^{2j+3l+6k}q^{\frac{j+k+l}{2}}\Lambda_j(\chi^2)\Lambda_l(\chi^3)Sym^k(\chi^6) \tag{4.19}$$

Hence the coefficient of u^n is indeed given by (4.15).

For $\chi\neq\chi_0,\chi_2,\chi_3$ mod Q for which at least one of χ^2,χ^3,χ^6 is not primitive or not odd, we still have $L(u,\chi)=\prod_{j=1}^{\deg Q-1}(1-\alpha_j(\chi)u)$ with all the inverse roots $|\alpha_j(\chi)|=q^{\frac{1}{2}}$ or $|\alpha_j(\chi)|=1$, and hence we obtain (4.16). □

Next, we want to evaluate $\sum_{\substack{2j+3l+6k=n\\0\leq j\leq N\\0\leq l\leq N\\0\leq k}}q^{j+k+l}$. Denote

$$S(n):=\sum_{\substack{2j+3l+6k=n\\0\leq j\leq N\\0\leq l\leq N\\0\leq k}}q^{j+k+l} \tag{4.20}$$

Lemma 4.2. *Let $N:=\deg Q-1$ then in the limit $q\to\infty$,*
for $0\leq n\leq 2N$

$$S(n)\sim q^{\lfloor\frac{n}{2}\rfloor} \tag{4.21}$$

for $2N<n\leq 5N$

$$S(n)\sim q^{\lfloor\frac{n+N}{3}\rfloor} \tag{4.22}$$

for $5N<n$

$$S(n)\sim q^{\frac{n+N}{6}}\cdot q^{\frac{-\lambda_n}{6}} \tag{4.23}$$

where

$$\lambda_n = \begin{cases} 0 & n = 5N \mod 6 \\ 7 & n = 5N+1 \mod 6 \\ 6 & n = 5N+2 \mod 6 \\ 3 & n = 5N+3 \mod 6 \\ 4 & n = 5N+4 \mod 6 \\ 11 & n = 5N+5 \mod 6 \end{cases} \tag{4.24}$$

Proof. In order to find the leading order term of $S(n)$ we need first to take the maximal possible j and then the maximal possible l which satisfy $2j + 3l + 6k = n, 0 \le j \le N, 0 \le l \le N$ (note that k will then be determined).

In the first case $0 \le n \le 2N$ if 2 divides n then $j = n/2, l = 0, k = 0$ will clearly give the leading order term. If 2 does not divide n then $j = (n-3)/2, l = 1, k = 0$ will give the leading order term.

In the second case $2N < n \le 5N$, we write $j = N - i_j$ and then we have $n - 2N = 6k + 3l - 2i_j$ and so clearly the values for k, l, i_j that will give the leading order term, depend on the value of $n - 2N \mod 3$ (or equivalently $n+N \mod 3$). Here our first priority is to minimize i_j and then to maximize l. The leading order term will be given by $q^{\frac{n+N-i_j}{3} - \frac{k}{2}}$ which gives $q^{\lfloor \frac{n+N}{3} \rfloor}$.

In the last case $5N < n$, we write $j = N - i_j$ and $l = N - i_l$, then we have $n - 5N = 6k - 3i_l - 2i_j$ ans so clearly the values for k, i_l, i_j that will give the leading order term, depend on the value of $n - 5N \mod 6$. Here our first priority is to minimize i_j and then to minimize i_l. The leading order term will be given by $q^{\frac{n+7N-3i_l-4i_j}{6}}$. Note that in the notations of (4.23) and (4.24) we have $4i_j + 3i_l = \lambda_n$

□

4.6. **Proof of Theorem 2.1.** Recall that by Lemma 3.2, we have that the number of characters which are not in $\Gamma_{d-prim}^{d-odd}(Q)$ is $O(\frac{\Phi(Q)}{q})$ when $q \to \infty$. Therefore, by using the formula for the variance that was given in (4.10) we may write

(4.25)

$$Var(S_{\alpha_2;n;Q}) = \frac{1}{\Phi(Q)^2} \sum_{\substack{\chi \neq \chi_0 \\ \chi \in \Gamma_{6-prim}^{6-odd}(Q)}} |\mathcal{M}(n;\alpha_2\chi)|^2 + \frac{1}{\Phi(Q)^2}|\mathcal{M}(n;\alpha_2\chi_2)|^2$$
$$+ \lambda_3 \frac{1}{\Phi(Q)^2}|\mathcal{M}(n;\alpha_2\chi_3)|^2 + O(\frac{S(n)}{\Phi(Q)q})$$

Where $\lambda_3 = 2$ if $|Q| = 1 \mod 3$, and zero otherwise. Note that we assume Q is prime and that the characteristic of the field $\mathbb{F}_q$ is odd therefor there is one quadratic character mod Q and either two or zero cubic characters, depending on if $|Q| = 1 \mod 3$ or not.

For the first sum, use equation (4.15) to have

$$\frac{1}{\Phi(Q)^2}\sum_{\substack{\chi\neq\chi_0\\ \chi\in\Gamma_{6-prim}^{6-odd}(Q)}}\Big|\sum_{\substack{2j+3l+6k=n\\ 0\leq j\leq N\\ 0\leq l\leq N\\ 0\leq k}} q^{\frac{j+k+l}{2}}\Lambda_j(\chi^2)\Lambda_l(\chi^3)Sym^k(\chi^6)\Big|^2 \tag{4.26}$$

We can use now the equidistribution result given in Theorem 3.8, to have

$$\frac{1}{\Phi(Q)}\iiint\limits_{U(N)}\Big|\sum_{\substack{2j+3l+6k=n\\ 0\leq j\leq N\\ 0\leq l\leq N\\ 0\leq k}} q^{\frac{j+k+l}{2}}\operatorname{tr}\Lambda_j(U_1)\operatorname{tr}\Lambda_l(U_2)\operatorname{tr}\mathrm{Sym}^k(U_3)\Big|^2\,dU_1\,dU_2\,dU_3 \tag{4.27}$$

It is well known that Λ_j and Sym^j are distinct irreducible representations of the unitary group $U(N)$, and hence one gets

$$\int\limits_{U(N)}\operatorname{tr}\Lambda_j(U)\overline{\operatorname{tr}\Lambda_i(U)}dU=\delta_{j,i} \tag{4.28}$$

and

$$\int\limits_{U(N)}\operatorname{tr}\mathrm{Sym}^j(U)\overline{\operatorname{tr}\mathrm{Sym}^i(U)}dU=\delta_{j,i} \tag{4.29}$$

Therefore

$$\frac{1}{\Phi(Q)^2}\sum_{\substack{\chi\neq\chi_0\\ \chi\in\Gamma_{6-prim}^{6-odd}(Q)}}|\mathcal{M}(n;\alpha_2\chi)|^2\sim\frac{1}{\Phi(Q)}\sum_{\substack{2j+3l+6k=n\\ 0\leq j\leq N\\ 0\leq l\leq N\\ 0\leq k}} q^{j+k+l} \tag{4.30}$$

The contribution from the second and third summands was evaluated in subsection 4.4. Adding up everything and checking for the leading order terms by using Lemma 4.2 finishes the proof.

5. The variance over short intervals

5.1. **The mean value.** The mean value of $\mathcal{N}_{\alpha_2;h}$ when we average over $A\in\mathcal{M}_n$ is

$$\begin{aligned}\langle\mathcal{N}_{\alpha_2;h}\rangle&:=\frac{1}{q^n}\sum_{A\in\mathcal{M}_n}\mathcal{N}_{\alpha_2;h}(A)\\ &=\frac{1}{q^n}\sum_{A\in\mathcal{M}_n}\sum_{f\in I(A;h)}\alpha_2(f)\\ &=q^{h+1}\frac{1}{q^n}\sum_{f\in\mathcal{M}_n}\alpha_2(f)\\ &=q^{h+1}\langle\alpha_2\rangle_n\end{aligned} \tag{5.1}$$

By (2.6) we have in the limit $q \to \infty$

$$\langle \mathcal{N}_{\alpha_2;h} \rangle \sim H \cdot q^{\lfloor n/2 \rfloor - n} \tag{5.2}$$

In the rest of this section we will evaluate the variance of $\mathcal{N}_{\alpha_2;h}$ i.e. the average of the squared difference between $\mathcal{N}_{\alpha_2;h}$ and its mean value.

5.2. **An expression for the variance.** To begin the proof of Theorem 2.2, we express the variance of the short interval sums $\mathcal{N}_{\alpha_2;h}$ in terms of sums of the function α_2, twisted by primitive even Dirichlet characters, similarly to what was done in the previous section.

Lemma 5.1. *As $q \to \infty$*

(5.3)

$$\mathrm{Var}(\mathcal{N}_{\alpha_2;h}) = \frac{H}{q^n} \frac{1}{\Phi^{ev}(T^{n-h})} \sum_{\substack{\chi \bmod T^{n-h} \\ \chi \neq \chi_0 \text{ even}}} \sum_{\substack{m_1,m_2=0 \\ m_1,m_2 \neq n-1}}^{n} \mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)}$$

where the definition of $\mathcal{M}(n;\alpha_2\chi)$ was first given in (4.8)

$$\mathcal{M}(n;\alpha_2\chi) := \sum_{f \in M_n} \alpha_2(f)\chi(f)\ . \tag{5.4}$$

Proof. To compute the variance, we use [14, Lemma 5.4] which gives an expression for the variance of sums over short intervals of certain arithmetic functions α which are "even" ($\alpha(cf) = \alpha(f)$ for $c \in \mathbb{F}_q^\times$), multiplicative, and symmetric under the map $f^*(t) := t^{\deg f} f(\frac{1}{t})$, in the sense that

$$\alpha(f^*) = \alpha(f), \quad \text{if } f(0) \neq 0\ . \tag{5.5}$$

Since the indicator function for square full polynomials α_2 clearly satisfies all of these conditions, we may use [14, Lemma 5.4] to obtain

$$\begin{aligned} \mathrm{Var}(\mathcal{N}_{\alpha_2;h}) = \frac{H}{q^n} &\sum_{m_1,m_2=0}^{n} \alpha_2(T^{n-m_1})\overline{\alpha_2(T^{n-m_2})} \\ &\times \frac{1}{\Phi^{ev}(T^{n-h})} \sum_{\substack{\chi \bmod T^{n-h} \\ \chi \neq \chi_0 \text{ even}}} \mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)} \end{aligned} \tag{5.6}$$

By the definition of α_2 we have $\alpha_2(T^{n-m}) = 1$ when $m \neq n-1$ and $\alpha_2(T^{n-m}) = 0$ when $m = n-1$, hence (5.3) follows. □

To compute the variance, we need to obtain an expression for $\mathcal{M}(n;\alpha_2\chi)$. Consider the generating function

$$\sum_{n=0}^{\infty} \mathcal{M}(n;\alpha_2\chi)u^n = \frac{L(u^2,\chi^2)L(u^3,\chi^3)}{L(u^6,\chi^6)} \tag{5.7}$$

For an even primitive characters χ mod T^{n-h}, $L(u,\chi)$ has a trivial zero at $u=1$, hence we may write

$$L(u,\chi)=(1-u)\det(I-uq^{\frac{1}{2}}\Theta_\chi)=(1-u)\sum_{i=0}^{n-h-2}u^iq^{i/2}\operatorname{tr}\Lambda_i(\Theta_\chi) \tag{5.8}$$

and

$$\frac{1}{L(u,\chi)}=\frac{1}{(1-u)\det(I-uq^{\frac{1}{2}}\Theta_\chi)}=\frac{1}{(1-u)}\sum_{i=0}^{\infty}u^iq^{i/2}\operatorname{tr}Sym^i(\Theta_\chi) \tag{5.9}$$

Therefore, the generating function of $\mathcal{M}(n;\alpha_2\chi)$ (i.e. $\frac{L(u^2,\chi^2)L(u^3,\chi^3)}{L(u^6,\chi^6)}$) for χ mod T^{n-h} such that χ^2,χ^3,χ^6 are primitive and even can be written as follows

$$\begin{aligned}&\frac{(1-u^2)(1-u^3)\det(I-u^2q^{\frac{1}{2}}\Theta_{\chi^2})\det(I-u^3q^{\frac{1}{2}}\Theta_{\chi^3})}{(1-u^6)\det(I-u^6q^{\frac{1}{2}}\Theta_{\chi^6})}=\\&\frac{(1-u^2)}{(1+u^3)}\sum_{j=0}^{n-h-2}\sum_{l=0}^{n-h-2}\sum_{k=0}^{\infty}u^{2j+3l+6k}q^{\frac{j+k+l}{2}}\Lambda_j(\chi^2)\Lambda_l(\chi^3)Sym^k(\chi^6)\end{aligned} \tag{5.10}$$

By expanding and comparing coefficients we have

$$\mathcal{M}(m;\alpha_2\chi)=\mathcal{S}'_\chi(m)-\mathcal{S}'_\chi(m-2) \tag{5.11}$$

where for $1\neq m\geq 0$

$$\mathcal{S}'_\chi(m):=\sum_{\substack{2j+3l+6k+3i=m\\0\leq j,l\leq n-h-2\\0\leq k,i}}(-1)^iq^{\frac{j+k+l}{2}}\Lambda_j(\chi^2)\Lambda_l(\chi^3)Sym^k(\chi^6) \tag{5.12}$$

and

$$\mathcal{S}'_\chi(1),\mathcal{S}'_\chi(-1),\mathcal{S}'_\chi(-2):=0. \tag{5.13}$$

Now back to the variance formula (see equation (5.3)), we can split the sum into two parts: the sum over $\chi\neq\chi_0 \mod T^{n-h},\chi\in\Gamma^{ev}_{prim}(T^{n-h})$ and the sum over even non-primitive characters mod T^{n-h}. We start by considering the first sum which will give the main term since most of the even characters are also primitive. With the second sum which will give an error term we deal later. Note: by Lemma 3.4 it is enough to split to these sums, and we can still use (5.9) and (5.8) for χ^2,χ^3,χ^6.

For χ even and primitive, consider the inner sum in the variance formula:

(5.14)
$$\sum_{\substack{m_1,m_2=0\\ m_1,m_2\neq n-1}}^{n} \mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)} = \sum_{m_1,m_2=0}^{n-2} \mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)} + |\mathcal{M}(n;\alpha_2\chi)|^2 + \overline{\mathcal{M}(n;\alpha_2\chi)}\sum_{m=0}^{n-2}\mathcal{M}(m;\alpha_2\chi) + \mathcal{M}(n;\alpha_2\chi)\sum_{m=0}^{n-2}\overline{\mathcal{M}(m;\alpha_2\chi)}$$

The sum over $\mathcal{M}(m;\alpha_2\chi)$ equals

(5.15)
$$\sum_{m=0}^{n-2}\mathcal{M}(m;\alpha_2\chi) = \sum_{m=0}^{n-2}({\mathcal{S}'}_{\chi}(m) - {\mathcal{S}'}_{\chi}(m-2)) = ({\mathcal{S}'}_{\chi}(n-2) + {\mathcal{S}'}_{\chi}(n-3))$$

Hence we get

(5.16)
$$\sum_{\substack{m_1,m_2=0\\ m_1,m_2\neq n-1}}^{n} \mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)} = |{\mathcal{S}'}_{\chi}(n)|^2 + |{\mathcal{S}'}_{\chi}(n-3)|^2 + {\mathcal{S}'}_{\chi}(n-3)\overline{{\mathcal{S}'}_{\chi}(n)} + \overline{{\mathcal{S}'}_{\chi}(n-3)}{\mathcal{S}'}_{\chi}(n)$$

By using the equidistribution result 3.6, the average of ${\mathcal{S}'}_{\chi}(m_1)\overline{{\mathcal{S}'}_{\chi}(m_1)}$ over all primitive even characters mod T^{n-h} in the limit $q\to\infty$ is

(5.17)
$$\frac{1}{\Phi^{ev}(T^{n-h})}\sum_{\substack{\chi \bmod T^{n-h}\\ \chi\neq\chi_0\in\Gamma^{ev}_{prim}}} {\mathcal{S}'}_{\Theta_\chi}(m_1)\overline{{\mathcal{S}'}_{\Theta_\chi}(m_2)} \sim$$
$$\iiint_{PU(n-h-2)} \sum_{\substack{2j+3l+6k+3i=m_1\\ 0\le j,l\le n-h-2\\ 0\le i,k}} (-1)^i q^{\frac{j+k+l}{2}} \operatorname{tr}\Lambda_j(U_1)\operatorname{tr}\Lambda_l(U_2)\operatorname{tr}\operatorname{Sym}^k(U_3)\times$$
$$\sum_{\substack{2j+3l+6k+3i=m_2\\ 0\le j,l\le n-h-2\\ 0\le i,k}} (-1)^i q^{\frac{j+k+l}{2}}\overline{\operatorname{tr}\Lambda_j(U_1)\operatorname{tr}\Lambda_l(U_2)\operatorname{tr}\operatorname{Sym}^k(U_3)}\,dU_1\,dU_2\,dU_3$$

By (4.28) and (4.29), we conclude that

(5.18)

$$\frac{1}{\Phi^{ev}(T^{n-h})}\sum_{\substack{m_1,m_2=0\\ m_1,m_2\neq n-1}}^{n}\mathcal{M}(m_1;\alpha_2\chi)\overline{\mathcal{M}(m_2;\alpha_2\chi)}=$$

$$\sum_{\substack{2j+3l+6k+3i=n\\ 0\leq j,l\leq n-h-2\\ 0\leq i,k}}q^{j+k+l}+\sum_{\substack{2j+3l+6k+3i=n-3\\ 0\leq j,l\leq n-h-2\\ 0\leq i,k}}q^{j+k+l}-2\sum_{\substack{2j+3l+6k+3i=n-3\\ 0\leq j,l\leq n-h-2\\ 0\leq i,k}}q^{j+k+l}=$$

$$\sum_{\substack{2j+3l+6k=n\\ 0\leq j,l\leq n-h-2\\ 0\leq k}}q^{j+k+l}$$

this combined with Lemma 4.2 (with $N=n-h-2$) gives the main term. Now, in order to complete the proof, it remains to bound the contribution of the even characters which are non-primitive to the variance. Note that we do not have quadratic or cubic characters here since $\Phi^{ev}(T^m)=q^{m-1}$ and we are considering the case of characteristic bigger than 13 (see 3.6), therefore there cannot be any even characters mod T^m of order 2 or 3. For even characters which are non-primitive we still have the bound (4.16) with $N=n-h-2$, and since their proportion is $O(1/q)$ in the set of even characters, then we can bound their contribution as in the previous section, therefore we skip the verification. Note: when bounding the contribution of non-primitive even characters we clearly don't use the equidistribution theorem as we do for the main term, therefore we cannot restrict to diagonal terms as in (5.18). However, this does not change the bound since the off-diagonal terms do not contribute higher order terms then the diagonal terms and their number does not depend on q.

Acknowledgements

The author gratefully acknowledges support under EPSRC Programme Grant EP/K034383/1 LMF: L-Functions and Modular Forms. The author would like to thank Zeev Rudnick for suggesting this problem and to both Jon Keating and Zeev Rudnick for helpful discussions and remarks.

References

[1] Bateman, Paul T.; Grosswald, Emil. *On a theorem of Erds and Szekeres.* Illinois J. Math. 2 (1958), no. 1, 88–98.

[2] Y. Cai, *On the distribution of square-full integers.* Acta Math. Sinica (N.S.) 13 (1997), 269280. A Chinese summary appears in Acta Math. Sinica 40 (1997),480.

[3] X.-D. Cao, *The distribution of square-full integers*, Period. Math. Hungar. 28 (1994), 4354.

[4] X. Cao, *On the distribution of square-full integers*, Period. Math. Hungar. 34 (1997), 169175.

[5] P. Erdös and S. Szekeres , *Über die Anzahl der Abelschen Gruppen gegebener Ordnung und über ein verwandtes zahlentheoretisches Problem*, Acta Univ.Szeged 7 (19341935), 95102.
[6] M. Filaseta and O. Trifonov. *The distribution of fractional parts with applications to gap results in number theory*. Proc. London Math. Soc. (3) 73 (1996), no. 2, 241278.
[7] D. R. Heath-Brown, *Square-full numbers in short intervals*. Square-full numbers in short intervals. Math. Proc. Cambridge Philos. Soc. 110 (1991), no. 1, 13.
[8] M. N. Huxley and O. Trifonov (1996). *The square-full numbers in an interval*. Mathematical Proceedings of the Cambridge Philosophical Society, 119, pp 201-208 doi:10.1017/S0305004100074107
[9] N. Katz. *Witt vectors and a question of Keating and Rudnick*. Int. Math. Res. Not. IMRN 2013, no. 16, 36133638.
[10] N. Katz. *On a question of Keating and Rudnick about primitive Dirichlet characters with squarefree conductor*. Int. Math. Res. Not. IMRN 2013, no. 14, 32213249.
[11] N. Katz. *Witt vectors and a question of Entin, Keating, and Rudnick*. Int. Math. Res. Not. IMRN 2015, no. 14, 59595975. doi: 10.1093/imrn/rnu120
[12] N. Katz. *On two questions of Entin, Keating, and Rudnick on primitive Dirichlet characters*. Int. Math. Res. Not. IMRN 2015, no. 15, 60446069. doi: 10.1093/imrn/rnu121
[13] J. P. Keating and Z. Rudnick. *The variance of the number of prime polynomials in short intervals and in residue classes*. Int. Math. Res. Notices, 2012; doi: 10.1093/imrn/rns220.
[14] J. P. Keating and Z. Rudnick. *Squarefree polynomials and Möbius values in short intervals and arithmetic progressions*. Algebra and Number Theory, vol 10., pp. 375-420
[15] H.-Q. Liu, *The distribution of square-full integers*, Ark. Mat. 32 (1994), 449454.
[16] H.-Q. Liu. *The number of squarefull numbers in an interval*. Acta Arith. 64 (1993), no. 2, 129149.
[17] H. Liu, and T. Zhang, *On the distribution of square-full numbers in arithmetic progressions*. Arch. Math. (Basel) 101 (2013), no. 1, 5364.
[18] M. Munsch, *Character sums over squarefree and squarefull numbers*. Arch. Math. (Basel) 102 (2014), no. 6, 555563.
[19] D. Suryanarayana and R. Sitamachandra, *The distribution of square-full integers*, Ark. Mat. (1973), 195201.
[20] J. Wu, *On the distribution of square-full integers*, Arch. Math. (Basel) 77 (2001), 233240.
[21] J. Wu, *On the distribution of square-full and cube-full integers*, Monatsh. Math. 126 (1998), 353367.
[22] W. Zhu and K. Yu, The distribution of square-full integers, Pure Appl. Math. (Xian) 12 (1996), 113122.

99

School of Mathematics, University of Bristol, Bristol BS8 1TW, UK
E-mail address: er14265@bristol.ac.uk